

Two-dimensional topological solitons in small exchange-dominated cylindrical ferromagnetic particles

Konstantin L. Metlov

Institute of Physics ASCR, Na Slovance 2, Prague 8, CZ-18221

(February 1, 2008)

A general approach allowing to construct the magnetization distributions of arbitrary topological charge in small exchange-dominated cylindrical ferromagnetic particles is presented. The exchange energy functional is minimized by these distributions exactly. The magnetostatic energy is accounted partially, so that it facilitates a choice between the topologically equivalent exchange-only solutions. The resulting magnetization distributions can be easily generalized to a variety of non-circular cylindrical shapes by means of a conformal transformation. As an example a magnetic structures of a thin circular ferromagnetic cylinder both with centered and displaced magnetic vortex and of a finite Bloch domain wall in an elongated particle is given.

PACS # 75.60.Ch, 75.70.Kw, 85.70.Kh

The magnetic structures of small ferromagnetic elements recently attained great experimental interest due to their possible applications in the integrated semiconductor-magnetic circuits and the magnetic random access memory (MRAM). From the point of view of these applications particularly the thin film elements (e.g. particles having cylindrical shape) are more interesting than others because they are more closely compatible with the current production technology. Currently, a lot of attention was devoted to modelling the magnetic structure of these elements [1–5] (see also references therein) with two main systematic approaches: the usage of guessed trial functions and the numerical solution of the Landau-Lifshitz equations. In this work an approximate (in accounting for magnetostatic interaction) but analytic approach for solving this task for a variety of element shapes is presented.

Mathematically, the problem of finding the static distribution of the magnetization vector $\vec{M}(\vec{R})$ ($\vec{R} = \{X, Y, Z\}$ denotes the radius vector of a point in space) within an isotropic ferromagnetic particle in zero external magnetic field can be formulated as finding the minimum of the following energy functional of $\vec{m}(\vec{R}) = \vec{M}(\vec{R})/M_S$:

$$M_S^2 e[\vec{m}] = \int_V \left\{ \frac{L_E^2}{2} \sum_{i=X,Y,Z} (\vec{\nabla} m_i)^2 - \vec{h}_D[\vec{m}] \cdot \vec{m} \right\} d^3 \vec{R}, \quad (1)$$

with the constraint $|\vec{m}(\vec{R})| = 1$ for $\vec{R} \in V$, where V denotes the volume of the particle, $L_E = \sqrt{C/M_S^2}$ is the exchange length, C is the exchange constant of the material, M_S is the saturation magnetization of the material, $\vec{\nabla} = \{\partial/\partial X, \partial/\partial Y, \partial/\partial Z\}$ is the gradient operator, $\vec{h}_D[\vec{m}] = \vec{H}_D[\vec{m}]/M_S$ is the demagnetizing field (a

functional of $\vec{m}$) created by the magnetization distribution. The demagnetizing field can be expressed with the help of Maxwell equations through its scalar potential $\vec{H}_D = \vec{\nabla} U(\vec{R})$, which is, in turn, a solution of the Poisson equation $\vec{\nabla}^2 U = \vec{\nabla} \cdot \vec{M}$ with the requirement (due to the finite size of the particle) that both $|\vec{R}|U$ and $|\vec{R}|^2|\nabla U|$ are finite as $|\vec{R}| \rightarrow \infty$, [6]. The main difficulty in treating this problem analytically arises from the fact that the dipolar interaction is non-local (as expressed by a functional dependence of $\vec{h}_D$ on $\vec{m}$), and the Euler-Lagrange condition for the extremum of the functional (1) yields a system of non-linear three-dimensional integral partial differential equations.

The simplified variant of the problem when the demagnetizing field is assumed to be zero $\vec{h}_D \equiv 0$ (so that the distribution is defined by a system of partial differential equations, but still non-linear ones) and V is taken as infinite two-dimensional space with the constraint $\vec{m}(\vec{R}) = m_0$ at $|\vec{R}| \rightarrow \infty$ was solved by Belavin and Polyakov [7,8]. Let us briefly rederive this solution because the procedure will be relevant further in this paper. Noting that the constraint $|\vec{m}| = 1$ leaves only two independent components of the magnetization vector, and parametrizing the magnetization vector by a complex function $w(z, \bar{z})$ of a complex variable $z = X + iY$, $\bar{z} = X - iY$, $i = \sqrt{-1}$ so that $m_x + im_y = 2w/(1 + w\bar{w})$ and $m_z = (1 - w\bar{w})/(1 + w\bar{w})$, the exchange energy functional can be written as

$$e = L_E^2 L_Z \int_V \frac{4}{(1 + w\bar{w})^2} \left(\frac{\partial w}{\partial z} \frac{\partial \bar{w}}{\partial \bar{z}} + \frac{\partial w}{\partial \bar{z}} \frac{\partial \bar{w}}{\partial z} \right) d^2 \vec{R}_2, \quad (2)$$

where $\partial/\partial z = (\partial/\partial X - i\partial/\partial Y)/2$, $\partial/\partial \bar{z} = (\partial/\partial X + i\partial/\partial Y)/2$ and $\vec{R}_2$ is a two-dimensional radius vector. The system of Euler equations for the extremum of this functional noting that the variables z and $\bar{z}$ are independent is

$$\frac{\partial}{\partial z} \left(\frac{\partial w}{\partial \bar{z}} \right) = \frac{2\bar{w}}{1 + w\bar{w}} \frac{\partial w}{\partial z} \frac{\partial w}{\partial \bar{z}}. \quad (3)$$

It is immediately clear that any analytic function w of the complex variable z satisfies this equation. Because the Cauchy-Riemann conditions for the analyticity $\partial w/\partial \bar{z} = 0$, if satisfied, turn both sides of (3) to zero. The Belavin-Polyakov (BP) soliton corresponds to the representation of $w(z)$ as a rational function of z . Zeroes of $w(z)$ correspond to the centers of solitons (at these points $m_z = 1$), at the infinity $|z| \rightarrow \infty$ the BP soliton has $m_z = -1$.

Let us now turn to small particles in the form of cylinders, choosing the coordinate system in such a way that the axis Z is perpendicular to the faces of the cylinder. As long as the thickness of these cylinders is much smaller than the exchange length it is reasonable to assume that the magnetization distribution is uniform along the thickness (Z axis). This assumption immediately reduces the dimensionality of the problem and allows us to use the $\vec{m}(\vec{r})$ parametrization in terms of $w(z, \bar{z})$.

A special case of the problem (1) taking into account the demagnetizing field was solved by Usov [9] for the particle shaped as a circular cylinder and is also recently confirmed experimentally [10]. This solution has the topology of a vortex, see Fig. . Its interesting property is that only a part of the magnetization distribution is a BP soliton (Usov calls this part the vortex core), the other part (represented in cylindrical coordinate system) as $m_r = 1$ is not. The component of magnetization normal to the cylinder face m_z changes from $m_z = 1$ at the center of the magnetic vortex to $m_z = 0$ at the vortex core boundary, and continues to stay zero in the non-core region.

The non-core part of the solution by Usov can be easily understood on the basis of the magnetic pole avoidance principle [6], which is due to the fact that the energy of the demagnetizing field $\vec{h}_D$ is always positive, and, since this field is created by magnetic poles (whose density is $\vec{\nabla} \cdot \vec{M}$), the system tries to avoid having poles. At a boundary of a finite magnetic particle where $|m|$ abruptly changes from 1 to 0 (no magnetic material) the density of magnetic poles is proportional to the component of the magnetization vector normal to the boundary. Thus, compared to the BP solution in the whole circular cylinder the solution by Usov has less magnetic poles on cylinder faces and is more favorable from the point of view of magnetostatic energy. Unfortunately, the procedure employed by Usov can hardly be generalized to other particle shapes because it heavily relies on the cylindrical coordinate system.

If we look back at equation (3) we can easily suspect that it may have more solutions besides the BP solitons. Although it was shown that the BP solitons are the only solutions with the finite total exchange energy in the infinite space, this requirement becomes irrelevant as long as we consider finite particles. Another family of solutions of (3) was first found by Gross [11] and is called merons:

they are expressed as $w_M(z, \bar{z}) = f(z)/\sqrt{f(z)\bar{f}(\bar{z})}$, where $f(z)$ is again an arbitrary analytic function. These solutions have $m_z = 0$ everywhere and do not produce magnetic poles on the cylinder faces. However, they diverge at the points where $f(z) = 0$ so that the exchange energy density is infinite at these points, therefore, the pure meron solutions can not be realized in the particles.

We may note, however, that on the contour $|f(z)| = 1$ the meron continuously joins the BP soliton $w(z) = f(z)$. Thus, the function

$$w_{SM}(z, \bar{z}) = \begin{cases} f(z) & |f(z)| \leq 1 \\ f(z)/\sqrt{f(z)\bar{f}(\bar{z})} & |f(z)| > 1 \end{cases} \quad (4)$$

is also a solution of (3) for any analytical function of complex variable $f(z)$, is regular and continuous everywhere and has much less magnetic poles than the BP solitons. This solution for the magnetization distributions of small magnetic particles is the main result of this paper. By choosing the appropriate $f(z)$ many realistic magnetization distributions can be obtained, some of the examples will be given later in the text.

The solution (4), generally, should have lower magnetostatic energy than the corresponding BP soliton $w(z) = f(z)$ because of less magnetic charges on the faces of the cylinder. However, such an advantage may turn out as unimportant for cylinders with a very small area of the region $f(z) > 1$. Unfortunately, the magnetostatic energy is strongly shape dependent due to its long-range character and the estimates of this critical particle size need to be done separately in each case. Once the energy of the solution (4) is lower, the BP soliton is unstable with respect to the continuous transformation to (4) by tilting spins in the region $|f(z)| > 1$ towards the plane of the cylinder. Such a transformation does not have an associated energy barrier because $w_{SM}(z)$ and $f(z)$ have the same topology.

Another important fact, easily checked by direct substitution, is that for any analytic function $o(z)$, the function $w(o(z), \bar{o}(\bar{z}))$ solves the equation (3) if $w(z, \bar{z})$ solves it. This property allows us to generalize the solution (4) using conformal transformations to particle shapes where the function $f(z)$ is hard to choose on the basis of symmetry argument alone.

As an illustration consider a particle shaped as a circular cylinder. From the symmetry it is natural to choose the BP solution in the form of a vortex centered in the cylinder (chosen as the origin of the coordinate system), which is $f(z) = \iota(z/c)^{|\nu|}$, where c is an arbitrary real number defining the scale and ν is the integer topological charge of the vortex. In the case $\nu = 1$ this solution is equivalent to the one obtained and investigated both analytically and numerically by Usov [9,12] and shown in Fig. . The definite value of c for a particular material and the geometry of the particle can be found by minimizing the total energy (1) with respect to c similarly as it is done in [9,12].

To displace the vortex the conformal transformation of the unit circle to the unit circle can be used, which is expressed by the rational function $o(z) = (z-a)/(1-\bar{a}z)$, where $a = X_c + iY_c$ is the new coordinate of the vortex center, $|a| < 1$. The resulting configuration (4) with $f(z) = \iota(z-a)/(1-\bar{a}z)/c$ is shown in Fig. . It can be seen from the picture that this configuration corresponds well (at least qualitatively) to the application of an external field in the direction of Y axis. Also the case of $|a| \rightarrow 1$ corresponds to the case of the uniformly magnetized particle. The complete analysis of the energies (both Zeeman and the magnetostatic) and the stability of such




FIG. 1. The structure of the vortex with $\nu = 1$ and $c = 1/2$ centered in the cylindrical particle. The outer solid circle is the contour of the particle, the inner circle is the boundary of the vortex core.




FIG. 2. The structure of the vortex with $\nu = 1$ and $c = 1/2$ displaced by $1/2$ with respect to the particle center. Note that the vortex core has the circular boundary but is otherwise asymmetrical.

a displaced vortex in an external magnetic field will be published elsewhere [13].

Another example is a multi-vortex solution of an arbitrary topological charge, which can be built by choosing $f(z)$ in (4) as a product of powers $(z - z_i)^{\nu_i}$, where z_i and ν_i are the coordinate of the center and the topological charge of i -th vortex, respectively. In particular, when there is an infinite number of $\nu_i = 1$ vortices sitting on a straight line (say $0 < z < 1$) the resulting magnetization distribution $f(z) = \exp(\int_0^1 \log(z - x) dx) = (z - 1)(z/(z - 1))^z/e$ can be classified as a finite Bloch domain wall with ends at points 0 and 1. The curves tangential to the vector $\vec{m}$ are of elongated (but not exactly elliptical) shape. A similar distribution was indeed observed in elongated magnetic dots [14] and interpreted as a two-domain configuration.

To summarize, a general approach allowing to build the magnetization distributions with an arbitrary topological charge in small exchange-dominated cylindrical magnetic particles of variety of shapes is presented. The produced magnetization distributions minimize the exchange functional exactly and partially account for the magnetostatic interaction. There are two factors limiting the presented consideration to small particles only. The first is connected with the fact that the magnetization distribution is assumed to be uniform along the cylinder thickness. The second is that the exchange energy of the proposed solution (4) within a circle of a given radius increases logarithmically with the radius. It means that the exchange energy diverges with increasing the particle size, thus other non-uniform configurations may become more energetically favorable in larger particles.

The presented approach should coexist well with existing numerical finite-element methods of micromagnetics as a way to specify the initial configuration with given topological properties, and to subject it to a final numerical refinement.

This work was supported in part by the Grant Agency of the Czech Republic under projects 202/99/P052 and 101/99/1662. I would like to thank Dr. Vladimír Kamberský, Dr. Ivan Tomáš, and Dr. Konstantin Guslienkov for reading the manuscript and many valuable discussions.

-
- [1] Y. Ishii and M. Sato, J. Appl. Phys. **65**, 3146 (1989).
 - [2] A. Aharoni, J. Appl. Phys. **68**, 255 (1990).
 - [3] N. A. Usov and S. E. Peschany, J. Magn. Magn. Mat. **110**, L1 (1992).
 - [4] Z. Guo and E. Della-Torre, Journal of Applied Physics **75**, 5710 (1994).
 - [5] S. T. Chui and V. N. Ryzhov, Phys. Rev. Lett. **78**, 2224 (1997).
 - [6] A. Aharoni, *Introduction to the theory of ferromagnetism*

- (Oxford University Press, Oxford, 1996), iSBN 0 19 851791 2.
- [7] A. A. Belavin and A. M. Polyakov, ZETP lett. **22**, 503 (1975), (in russian).
 - [8] A. M. Kosevich, B. A. Ivanov, and A. S. Kovalev, Phys. Rep. **194**, 117 (1990).
 - [9] N. A. Usov and S. E. Peschany, J. Magn. Magn. Mat. **118**, L290 (1993).
 - [10] T. Shinjo *et al.*, Science **289**, 930 (2000).
 - [11] D. J. Gross, Nucl. Phys. B **132**, 439 (1978).
 - [12] N. A. Usov and S. E. Peschany, Fiz. Met. Metal **12**, 13 (1994), (in russian).
 - [13] K. Y. Guslienko and K. L. Metlov, Evolution and stability of a magnetic vortex in small cylindrical ferromagnetic particle under applied field. The reference to xxx.lanl.gov will be here (unpublished).
 - [14] J. Wittborn, K. V. Rao, J. Nogués, and I. K. Schuller, Appl. Phys. Lett. **76**, 2931 (2000).